

M/IV Order Sheet

Internal use only

Consideration should be given to patient specific customization

Date

INTRAVENOUS THERAPIES

☐ **Standard** Infuse over 1 hour
0.9% NSS 500 cc /week X
Ascorbic Acid 10000 mg
Magnesium Chloride 2000 mg
Calcium Gluconate 300 mg
B complex 200 mg
Dexpanthenol (B5) 250 mg
B6 300 mg
Lidocaine (2%) 140 mg
Na Bicarb 8.4% (8) cc
Acetyl-L-Carnitine 200 mg
B12 (Methylcobalamin) 3000 mcg
AMP 50 mg
Tyrosine 10 mg
D-ribose 15 mg
NADH 20 mg

Plus IV QUICK DRIP: Infuse over 10 mins following IV
Glutathione 1500 mg
Na Bicarb 8.4% 2 cc

☒ **Back & Muscle Pain** Infuse over 1 hour
Standard /week X
0.9% NSS 500 cc
Lidocaine 2% 100mg
Colchicine (IVP)** 1mg
**give colchicine IVP after infusion for 2-5 minutes

Plus IV QUICK DRIP: Infuse over 10 mins following IV
Glutathione 1500 mg
Na Bicarb 8.4% 2 cc

☐ **Viral Plus - INITIAL** Infuse over 1 hour
Standard /week X
0.9% NSS 500 cc
Thymus 20 mg
Glycyrhizinate 40mg
Ascorbic Acid 20000 mg

Plus IV QUICK DRIP: Infuse over 10 mins following IV
Glutathione 1500 mg
Na Bicarb 8.4% 2 cc

☐ **Viral Plus - SUBSEQUENT** Infuse over 1 hour
Standard /week X
0.9% NSS 500 cc
Thymus 20 mg
Glycyrhizinate 40mg
Ascorbic Acid 30000 mg

Plus IV QUICK DRIP: Infuse over 10 mins following IV
Glutathione 1500 mg
Na Bicarb 8.4% 2 cc

☐ **Memory** Infuse over 1 hour
D5W 250 cc /week X
Acetyl-L-Carnitine 400 mg
B12 (Methylcobalamin) 3000 mcg
AMP 50 mg
Tyrosine 10 mg
D-ribose 15 mg
Ginkgo 50 mg
Vinpocetine 10 mg
DMAE 300 mg
NADH 20 mg
Piracetam 400 mg

Plus IV QUICK DRIP: Infuse over 10 mins following IV
Glutathione 1500 mg
Na Bicarb 8.4% 2 cc

INTRAVENOUS SUPER VIRAL THERAPY

☐ **Super Viral** Infuse over 1 hour
0.45% NSS 500 cc /week X
Glycyrhizinate 400 mg
Thymus 20 mg
Ascorbic Acid 40000 mg
Na Bicarb 8.4% (8) cc

INTRAMUSCULAR INJECTIONS

☒ **Energy Injection** slow gluteal IM administration Freq
AMP 100 mg/cc (1cc)
NADH 10 mg/cc (1cc)
Glutathione 100 mg/cc (16 cc)
Hydroxycobalamin 1000mg/cc (1cc)
Na Bicarb 8.4% (0.4 cc)

☐ **Testosterone** mg Freq

☐ **Nandrolone** mg Freq

☐ **Testosterone** mg/Nandrolone mg Freq

☐ **Immunoglobulin** cc Freq

☐ **Other** Dose cc Freq

TRIGGER POINT INJECTIONS

Formula 1:
Kenalog 40 mg/cc (1.5 cc)
Hydroxocobalamin 1000 mcg/cc (3 cc or 4 cc)
Procaine 2% (2 cc)
Marcaine 0.25% (2 cc)
Depo Medrol 80 mg/cc (0.5 cc) - Optional ingredient

Formula 2:
1% Lidocaine without epinephrine

NURSING DOCUMENTATION

Pre-Infusion Vital Signs T 98.7 P 82 R 16 B/P 141/79

Post Infusion Vital Signs T 97.3 P 73 R 14 B/P 152/87

22 g angiocath inserted Site LAC
225 IV start time IV end time 5:30
578 Warm pack applied Cool pack applied
cc infused
IV catheter intact upon removal
IV site unremarkable following infusion/removal of catheter
Patient tolerated procedure well
Patient verbalized understanding of procedure
Adverse reaction/event (see documentation below)

Notes:

IM injection given Site of Administration:
Injection site unremarkable following procedure
Patient tolerated procedure well
Patient verbalized understanding of procedure
Warm pack applied Cool pack applied

Notes:

IV SERIES

Nursing documentation for each infusion on additional IV/IM order sheet

Standard
Back/Muscle
Viral Plus
Memory
Super Viral
Energy Inj.
Testosterone
Nandrolone
Test/Nand
IMIG
Other

Enter Date Administered

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16

IV/IM Consents

IV Obtained
IM Obtained

Form No
FFCIV1 042507

Physician Signature

Nurse/MA Signature

Date